
11.8 Calculate: Angular momentum and kinetic energy of a 3D rotating rigid body.

Consider a rigid body B rotating in a Newtonian reference frame N. Right-handed orthogonal unit vectors $\hat{\mathbf{b}}_{\mathbf{x}}$, $\hat{\mathbf{b}}_{\mathbf{y}}$, $\hat{\mathbf{b}}_{\mathbf{z}}$ are fixed in B and parallel to B's principal inertia axes about $B_{\rm cm}$ (B's center of mass).



Quantity	Symbol	Type
B 's moment of inertia about $B_{\rm cm}$ for $\hat{\mathbf{b}}_{\rm x}$	I_{xx}	Constant
B 's moment of inertia about $B_{\rm cm}$ for $\hat{\mathbf{b}}_{\rm y}$	I_{yy}	Constant
B 's moment of inertia about $B_{\rm cm}$ for $\hat{\mathbf{b}}_{\rm z}$	I_{zz}	Constant
$[\hat{\mathbf{b}}_{x}, \hat{\mathbf{b}}_{y}, \hat{\mathbf{b}}_{z}]$ measures of ${}^{N}\vec{\boldsymbol{\omega}}^{B}$	$\omega_x, \omega_y, \omega_z$	Variables

$$\overset{N}{\boldsymbol{\omega}}^{B} = \omega_{x} \, \hat{\mathbf{b}}_{x} + \omega_{y} \, \hat{\mathbf{b}}_{y} + \omega_{z} \, \hat{\mathbf{b}}_{z}$$

$$\overset{\exists}{\mathbf{I}}^{B/B_{cm}} = \mathbf{I}_{xx} \, \hat{\mathbf{b}}_{x} \, \hat{\mathbf{b}}_{x} + \mathbf{I}_{yy} \, \hat{\mathbf{b}}_{y} \, \hat{\mathbf{b}}_{y} + \mathbf{I}_{zz} \, \hat{\mathbf{b}}_{z} \, \hat{\mathbf{b}}_{z}$$

$$= \begin{bmatrix} \mathbf{I}_{xx} & 0 & 0 \\ 0 & \mathbf{I}_{yy} & 0 \\ 0 & 0 & \mathbf{I}_{zz} \end{bmatrix}_{\hat{\mathbf{b}}_{xyz}}$$

Calculate
$$B$$
's angular momentum about B_{cm} in N and B 's rotational kinetic energy in N .

Result: ${}^{N}\vec{\mathrm{H}}^{B/B_{\text{cm}}} = \prod_{(15.1)} \mathrm{I}_{\mathrm{xx}} \, \omega_{x} \, \hat{\mathbf{b}}_{\mathrm{x}} + \mathrm{I}_{\mathrm{yy}} \, \omega_{y} \, \hat{\mathbf{b}}_{\mathrm{y}} + \mathbf{\hat{b}}_{\mathrm{z}}$
 $\hat{\mathbf{b}}_{\mathrm{z}}$
 ${}^{N}K^{B}_{\text{rotation}} = \frac{1}{2} \left(\mathrm{I}_{\mathrm{xx}} \, \omega_{x}^{2} + \mathrm{I}_{\mathrm{yy}} \, \omega_{y}^{2} + \mathbf{\hat{b}}_{\mathrm{z}} \right)$

11.9 3D spinning rigid body: Guess and check conservation of energy/angular momentum.


This problem refers to Homeworks 6.20 and 11.8.

$$\vec{M} = \begin{bmatrix} N_d \vec{H} \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \dot{\omega}_x = \begin{bmatrix} (I_{yy} - I_{zz}) \omega_z \omega_y \end{bmatrix} / I_{xx}$$

$$\Rightarrow \dot{\omega}_y = \begin{bmatrix} (I_{zz} - I_{xx}) \omega_x \omega_z \end{bmatrix} / I_{yy}$$

$$\dot{\omega}_z = \begin{bmatrix} (I_{xx} - I_{yy}) \omega_y \omega_x \end{bmatrix} / I_{zz}$$







- Using your intuition and/or conservation of angular momentum/mechanical energy (Sections 20.9 and 23.2), circle the quantities below that you guess remain constant (are "conserved").
- Next, solve the ODEs in Hw 6.20 for $0 \le t \le 4$ with initial values $\omega_x = 7$, $\omega_y = 0.2$, $\omega_z = 0.2$ (use

$$H_x = \vec{\mathbf{H}} \cdot \hat{\mathbf{b}}_x$$
 $H_y = \vec{\mathbf{H}} \cdot \hat{\mathbf{b}}_y$ $H_z = \vec{\mathbf{H}} \cdot \hat{\mathbf{b}}_z$ $|\vec{\mathbf{H}}|$

Hint: To determine if a quantity such as ω_x is constant during numerical integration, look at the numerical values of ω_x or **zoom** into the plot of ω_x vs. t and check if ω_x is constant to within a reasonable multiplier (e.g., 10) of numerical integrator accuracy. In MotionGenesis and MATLAB[®], the default numerical integrator accuracy is $\approx 1 \times 10^{-6}$.

Solution at www.MotionGenesis.com \Rightarrow Get Started \Rightarrow Solving 1st-order ODEs. Add the line: Hx = Ixx*wx; Hy = Iyy*wy; Hz = Izz*wz; Hmag = sqrt(Hx^2 + Hy^2 + Hz^2), etc. Add the line: Output t sec, Hx kg*m^2/sec, Hy kg*m^2/sec, Hz kg*m^2/sec, Hmag kg*m^2/sec, etc. View/plot the numerical results (numbers) and determine if they stay constant to within $\approx 1 \times 10^{-5}$.

Optional: Simulate delayed 3D spin instability for wingnut or T-handle with: $I_{xx} < I_{yy} < I_{zz}$, spin is initially mostly about $\hat{\mathbf{b}}_{\mathrm{y}}$ (intermediate moment-of-inertia axis), and $(I_{\mathrm{yy}}-I_{\mathrm{xx}})(I_{\mathrm{zz}}-I_{\mathrm{yy}})$ is small, e.g., by using $\omega_x(0)=0.2\,\frac{\mathrm{rad}}{\mathrm{sec}},\,\omega_y(0)=7.0\,\frac{\mathrm{rad}}{\mathrm{sec}},\,\omega_z(0)=0.2\,\frac{\mathrm{rad}}{\mathrm{sec}},\,I_{\mathrm{xx}}=1.9\,\mathrm{kg}\,\mathrm{m}^2,\,I_{\mathrm{yy}}=2.0\,\mathrm{kg}\,\mathrm{m}^2,\,I_{\mathrm{zz}}=3.0\,\mathrm{kg}\,\mathrm{m}^2,$ Video and Dzhanibekov analysis at <u>www.MotionGenesis.com</u> \Rightarrow <u>Get Started</u> \Rightarrow Spin stability





